

Magnesium/Potassium Aspartate

DESCRIPTION

Magnesium and potassium are minerals involved in important physiological functions that contribute to overall health.[‡]

INDICATIONS

- Support for heart, muscular and nerve health[‡]

FUNCTIONS AND MECHANISM OF ACTION

Magnesium and potassium provide key roles in heart, muscular and nerve health to support a number of physiological functions, including cardiovascular health and muscle function. Both minerals are essential for healthy nerve impulse function, muscle contraction and acid/alkaline balance. Magnesium supports cardiovascular health by promoting arterial and endothelial function, while potassium plays a critical role in the transmission of electrical impulses in the heart and supports healthy blood flow. Magnesium also is involved in carbohydrate and nutrient metabolism as it is responsible for activating the enzymes necessary for various physiological functions, including the metabolism of macronutrients, energy production and the utilization of calcium, phosphorus, sodium and potassium. In addition, magnesium plays an important role in facilitating potassium utilization and promotes healthy bone metabolism as it is an essential bone matrix mineral. Potassium supports bone cell function and bone mineralization in part by enhancing calcium absorption and supporting healthy bone resorption.[‡]

FORMULA (#202752)

Serving Size 1 Vegetarian Capsule

Magnesium (as magnesium aspartate)	70 mg
Potassium (as potassium aspartate)	99 mg

Other Ingredients: Hydroxypropyl methylcellulose (capsule), ascorbyl palmitate

Gluten-free, Non-GMO

SUGGESTED USE

As a dietary supplement, adults take 1 capsule, 1-4 times or as directed by your health professional.

WARNING

Not to be taken if you have hyperkalemia or abnormal kidney function, or take ACE inhibitors or potassium sparing diuretics. If you are pregnant or lactating, have any health condition or are taking any medication, consult your health professional before use.

Keep out of reach of children.

STORAGE

Store in a cool, dry place, away from direct light. Use only if safety seal is intact.

REFERENCES

- Dahle LO, Berg G, Hammar M, et al. *Am J Obstet Gynecol*. 1995 Jul;173(1):175-80.
Veronese N, Berton L, Carraro S, et al. *Am J Clin Nutr*. 2014 Sep;100(3):974-81.
Dorup I, Skajaa K, Clausen T, et al. *Br Med J (Clin Res Ed)*. 1988 Feb 13; 296(6620): 455–458.
Walker AF, De Souza MC, Vickers MF, et al. *J Womens Health*. 1998 Nov;7(9):1157-65.
Facchinetto F, Borella P, Sances G, et al. *Obstet Gynecol*. 1991 Aug;78(2):177-81.

Taubert K. *Fortschr Med.* 1994 Aug 30;112(24):328-30.
Grazzi L, Andrasik F, Usai S, et al. *Neurol Sci.* 2007 Jun;28(3):148-50.
Morskowitz MA. *Headache.* 2007. 47(1); S58-S63.
Minich DM, Bland JS. *Altern Ther Health Med.* 2007 Jul-Aug;13(4):62-5.
Matuszkiewicz-Rowińska J, Wojtaszek E. *Wiad Lek.* 2013;66(4):286-9.
Ji B, Liu J, Liu M, et al. *J Cardiovasc Surg (Torino).* 2006 Dec;47(6):671-5.
Singh RB, Singh NK, Niaz MA, Sharma JP. *Int J Clin Pharmacol Ther.* 1996 May;34(5):219-25.
Fuentes JC, Salmon AA, Silver MA. *Congest Heart Fail.* 2006 Jan-Feb;12(1):9-13.
Wirell MP, Wester PO, Stegmayr BG. *J Intern Med.* 1994 Aug;236(2):189-95.
Witteman JC, Grobbee DE, Derkx FH, et al. *Am J Clin Nutr.* 1994 Jul;60(1):129-35.
Mathew R, Gloster ES, Altura BT, Altura BM. *Clin Sci.* 1988;75, 661-667.
Pu J, Zhang C, Quan X, et al. *J Huazhong Univ Sci Technolog Med Sci.* 2008 Oct;28(5):517-9.
Franzoni F, Santoro G, Carpi A, et al. *Biomed Pharmacother.* 2005 Jan-Feb;59(1-2):25-9.
Braschi A, Naismith DJ. *Br J Nutr.* 2008 Jun;99(6):1284-92.
Naismith DJ, Braschi A. *Br J Nutr.* 2003 Jul;90(1):53-60.
Jee SH, Miller ER 3rd, Guallar E, et al. *Am J Hypertens.* 2002 Aug;15(8):691-6.
He K, Liu K, Daviglus ML, et al. *Circulation.* 2006 Apr 4;113(13):1675-82.
Mooren FC. *Diabetes Obes Metab.* 2015 Sep;17(9):813-23.
Nemoto T, Matsuzaki H, Uehara M, Suzuki K. *Magnes Res.* 2006 Mar;19(1):19-27.
Brilla LR, Haley TF. *J Am Coll Nutr.* 1992 Jun;11(3):326-9.
Toba Y, Kajita Y, Masuyama R, et al. *J. Nutr.* 2000. 130(2); 216-220.
Hiroshi M, Tomoko N, Shin-ichi K, et al. *Jpn J Nutr Diet.* 2005. 63(1); 27-31.
Hamill-Ruth RJ, McGory R. *Crit Care Med.* 1996 Jan;24(1):38-45.
Fordtran JS, Morawski SG, Santa Ana CA. *Gastroenterol.* 1985 Nov;89(5):1050-3.
Orchard TS, Larson JC, Alghothani N, et al. *Am J Clin Nutr.* 2014 Apr; 99(4): 926–933.
Dimai HP, Porta S, Wirnsberger G, et al. *J Clin Endocrinol Metab.* 1998 Aug;83(8):2742-8.
Lambert H, Frassetto L, Moore JB, et al. *Osteoporos Int.* 2015 Apr;26(4):1311-8.
Moseley KF, Weaver CM, Appel L, et al. *J Bone Miner Res.* 2013 Mar;28(3):497-504.
Buehlmeier J, Frings-Meuthen P, Remer T, et al. *J Clin Endocrinol Metab.* 2012 Dec;97(12):4789-97.
Ryder KM, Shorr RI, Bush AJ, et al. *J Am Geriatr Soc.* 2005 Nov;53(11):1875-80.
Jehle S, Zanetti A, Muser J, et al. *J Am Soc Nephrol.* 2006 Nov;17(11):3213-22.

For more information on Magnesium/Potassium Aspartate, visit douglaslabs.com

[†]These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by
Douglas Laboratories
600 Boyce Rd
Pittsburgh, PA 15205
800-245-4440
douglaslabs.com



PUSH YOUR POTENTIAL.

©2022 Douglas Laboratories. All Rights Reserved.