

DIM Enhanced Delivery System Promotes Healthy Estrogen*

DIM (diindolylmethane) has been shown to help regulate and promote a more efficient metabolism of estrogen, and an optimal ratio of estrogen metabolites.*

DIM Enhanced Delivery System contains BioResponse DIM®, a unique formulation containing pure diindolylmethane, an indole. Indoles are plant compounds with health promoting properties, and are found in cruciferous vegetables such as broccoli, cabbage, cauliflower and Brussels sprouts.* The phytochemicals in cruciferous vegetables have been shown to beneficially affect the body's hormonal and detoxification systems, and epidemiological studies support the health benefits of consuming these vegetables.*

Key Features

- Promotes healthy estrogen metabolism and balance*
- Helps promote the conversion of estrogen to its beneficial, protective 2-hydroxyestrone metabolites and reduces production of genotoxic 16α-hydroxyestrone*
- Stimulates detoxification enzyme systems*



Item #74140 120 vegetarian capsules





DIM Enhanced Delivery System

DIM is a major active acid-catalyzed derivative of one of the phytochemicals in cruciferous vegetables, indole-3-carbinol (I3C). DIM is thought to be responsible for the health effects of dietary I3C.* DIM Enhanced Delivery System is a stable, bioavailable form of DIM, made possible through a proprietary delivery system. The formula is co-solubilized with phosphatidylcholine, and microencapsulated in starch particles.

Research over the past thirty years has determined that healthy estrogen metabolism is closely linked to several healthy parameters in men and women, particularly some involving the breast, uterus, prostate and other reproductive tissue.* Genetics, excess weight, poor diet and other lifestyle factors may result in an imbalance of estrogen metabolites. Xenoestrogenic compounds, such as organochlorine pesticides, can also significantly disrupt healthy estrogen metabolism.

These estrogen disruptors alter estradiol hydroxylation metabolism producing a higher ratio of the genotoxic 16α -hydroxyestrone ($16\square$ -OHE1) to the safer and weaker estrogenic 2-hydroxyestrone (2-OHE1).* The genotoxic 16α -OHE1 can potentially disrupt several normal cellular metabolic processes.* DIM promotes the conversion of estrogen to its beneficial, protective 2-hydroxyestrone metabolites and reduces production of genotoxic 16α -OHE1.* Modulating these aspects of estrogen metabolism, particularly the production of 16α -OHE1, may contribute to healthy aging.*

Supplement Facts	
Serving Size	4 Capsules
Servings Per Container	30
Amount Per Serving	% Daily Value
BioResponse DIM®	300 mg †
BioResponse DIM® (A proprietary Diindolylmethane complex - svitamin E (as tocophersolan), phosphatidyl	starch, DIM (25% min.).

Other ingredients: Hydroxypropyl methylcellulose, microcrystalline cellulose, rice hull concentrate, gum arabic, sunflower oil.

Suggested Use: As a dietary supplement, 2 to 4 capsules one or two times daily with food, or as directed by a healthcare practitioner.

Caution: Do not use this product if you are pregnant or lactating, or using birth control pills. Persons taking prescription medications should consult a healthcare professional before use. Harmless changes in urine color may occur. Increased water consumption reverses this side effect. Keep out of reach of children.

BioResponse DIM® is a proprietary, enhanced bioavailability complex containing diindolylmethane licensed from BioResponse, L.L.C., Boulder, Colorado.*

The mechanisms for DIM's health benefits primarily involve the induction of mixed function oxidases and phase II detoxification enzyme systems by the binding and activation of the arylhydrocarbon receptor (AhR).* Research using human breast cells (MCF-7) has shown that the binding of DIM® to the arylhydrocarbon receptor can result in rapid formation of the nuclear AhR complex and consequent induction of gene expression and synthesis of cytochrome P450 detoxification enzyme (CyP450AI).* DIM® consequently produces increased levels of the protective hydroxylated estrogen 2-OHEI.* Some have suggested that DIM may also positively affect cellular signaling pathways.*

It is becoming increasingly apparent that DIM may provide an important mechanism for supporting successful aging despite the increasing levels of xenoestrogenic compounds in our modern world.*

References:

Bell MC, Crowley-Nowick P, Bradlow HL, et al. Gynecol Oncol 2000;78:123-9. Ben-Jonathan N, Cooper RL, Foster P, et al. Environ Health Perspect 1999;107 Suppl 4:605-11.

Bradlow HL, Davis D, Sepkovic DW, et al. Sci Total Environ 1997;208:9-14.
Bradlow HL, Sepkovic DW, Telang NT, et al. Ann N Y Acad Sci 1999;889:204-13.
Chang YC, Riby J, Chang GH, et al. Biochem Pharmacol 1999;58:825-34. Chen I, Safe S, Bjeldanes L. Biochem Pharmacol 1996;51:1069-76.

Davis DL, Telang NT, Osborne MP, et al. Environ Health Perspect 1997;105 Suppl 3:571-6. Ge X, Fares FA, Yannai S. Anticancer Res 1999;19:3199-203. Lake BG, Tredger JM, Renwick AB, et al. Xenobiotica

Michnovicz JJ, Adlercreutz H, Bradlow HL. J Natl Cancer Inst 1997;89:718-23.

Riby JE, Chang GH, Firestone GL, et al. Biochem Pharmacol 2000:60:167-77.

Sanderson JT, Slobbe L, Lansbergen GW, et al. Toxicol Sci 2001;61:40-8. Shertzer HG, Senft AP. Drug Metabol Drug Interact 2000;17:159-88. Wattenberg LW, Loub WD. Cancer Res 1978;38:1410-3.